

## 1997 Calculus AB Solutions: Part A

1. C  $\int_1^2 (4x^3 - 6x) dx = (x^4 - 3x^2) \Big|_1^2 = (16 - 12) - (1 - 3) = 6$
  
2. A  $f(x) = x(2x - 3)^{\frac{1}{2}}; f'(x) = (2x - 3)^{\frac{1}{2}} + x(2x - 3)^{-\frac{1}{2}} = (2x - 3)^{-\frac{1}{2}}(3x - 3) = \frac{(3x - 3)}{\sqrt{2x - 3}}$
  
3. C  $\int_a^b (f(x) + 5) dx = \int_a^b f(x) dx + 5 \int_a^b 1 dx = a + 2b + 5(b - a) = 7b - 4a$
  
4. D  $f(x) = -x^3 + x + \frac{1}{x}; f'(x) = -3x^2 + 1 - \frac{1}{x^2}; f'(-1) = -3(-1)^2 + 1 - \frac{1}{(-1)^2} = -3 + 1 - 1 = -3$
  
5. E  $y = 3x^4 - 16x^3 + 24x^2 + 48; y' = 12x^3 - 48x^2 + 48x; y'' = 36x^2 - 96x + 48 = 12(3x - 2)(x - 2)$   
 $y'' < 0$  for  $\frac{2}{3} < x < 2$ , therefore the graph is concave down for  $\frac{2}{3} < x < 2$
  
6. C  $\frac{1}{2} \int e^{\frac{t}{2}} dt = e^{\frac{t}{2}} + C$
  
7. D  $\frac{d}{dx} \cos^2(x^3) = 2 \cos(x^3) \left( \frac{d}{dx} (\cos(x^3)) \right) = 2 \cos(x^3) (-\sin(x^3)) \left( \frac{d}{dx} (x^3) \right)$   
 $= 2 \cos(x^3) (-\sin(x^3)) (3x^2)$
  
8. C The bug change direction when  $v$  changes sign. This happens at  $t = 6$ .
  
9. B Let  $A_1$  be the area between the graph and  $t$ -axis for  $0 \leq t \leq 6$ , and let  $A_2$  be the area between the graph and the  $t$ -axis for  $6 \leq t \leq 8$ . Then  $A_1 = 12$  and  $A_2 = 1$ . The total distance is  $A_1 + A_2 = 13$ .
  
10. E  $y = \cos(2x); y' = -2 \sin(2x); y' \left( \frac{\pi}{4} \right) = -2$  and  $y \left( \frac{\pi}{4} \right) = 0; y = -2 \left( x - \frac{\pi}{4} \right)$
  
11. E Since  $f'$  is positive for  $-2 < x < 2$  and negative for  $x < -2$  and for  $x > 2$ , we are looking for a graph that is increasing for  $-2 < x < 2$  and decreasing otherwise. Only option E.
  
12. B  $y = \frac{1}{2}x^2; y' = x; \text{ We want } y' = \frac{1}{2} \Rightarrow x = \frac{1}{2}. \text{ So the point is } \left( \frac{1}{2}, \frac{1}{8} \right).$

## 1997 Calculus AB Solutions: Part A

13. A  $f'(x) = \frac{|4-x^2|}{x-2}$ ;  $f$  is decreasing when  $f' < 0$ . Since the numerator is non-negative, this is only when the denominator is negative. Only when  $x < 2$ .
14. C  $f(x) \approx L(x) = 2 + 5(x-3)$ ;  $L(x) = 0$  if  $0 = 5x - 13 \Rightarrow x = 2.6$
15. B Statement B is true because  $\lim_{x \rightarrow a^-} f(x) = 2 = \lim_{x \rightarrow a^+} f(x)$ . Also,  $\lim_{x \rightarrow b} f(x)$  does not exist because the left- and right-sided limits are not equal, so neither (A), (C), nor (D) are true.
16. D The area of the region is given by  $\int_{-2}^2 (5 - (x^2 + 1)) dx = 2 \left( 4x - \frac{1}{3} x^3 \right) \Big|_0^2 = 2 \left( 8 - \frac{8}{3} \right) = \frac{32}{3}$
17. A  $x^2 + y^2 = 25$ ;  $2x + 2y \cdot y' = 0$ ;  $x + y \cdot y' = 0$ ;  $y'(4, 3) = -\frac{4}{3}$ ;  
 $x + y \cdot y' = 0 \Rightarrow 1 + y \cdot y'' + y' \cdot y' = 0$ ;  $1 + (3)y'' + \left(-\frac{4}{3}\right) \cdot \left(-\frac{4}{3}\right) = 0$ ;  $y'' = -\frac{25}{27}$
18. C  $\int_0^{\frac{\pi}{4}} \frac{e^{\tan x}}{\cos^2 x} dx$  is of the form  $\int e^u du$  where  $u = \tan x$ .  
 $\int_0^{\frac{\pi}{4}} \frac{e^{\tan x}}{\cos^2 x} dx = e^{\tan x} \Big|_0^{\frac{\pi}{4}} = e^1 - e^0 = e - 1$
19. D  $f(x) = \ln|x^2 - 1|$ ;  $f'(x) = \frac{1}{x^2 - 1} \cdot \frac{d}{dx}(x^2 - 1) = \frac{2x}{x^2 - 1}$
20. E  $\frac{1}{8} \int_{-3}^5 \cos x dx = \frac{1}{8} (\sin 5 - \sin(-3)) = \frac{1}{8} (\sin 5 + \sin 3)$ ; Note: Since the sine is an odd function,  $\sin(-3) = -\sin(3)$ .
21. E  $\lim_{x \rightarrow 1} \frac{x}{\ln x}$  is nonexistent since  $\lim_{x \rightarrow 1} \ln x = 0$  and  $\lim_{x \rightarrow 1} x \neq 0$ .
22. D  $f(x) = (x^2 - 3)e^{-x}$ ;  $f'(x) = e^{-x}(-x^2 + 2x + 3) = -e^{-x}(x-3)(x+1)$ ;  $f'(x) > 0$  for  $-1 < x < 3$
23. A Disks where  $r = x$ .  $V = \pi \int_0^2 x^2 dy = \pi \int_0^2 y^4 dy = \frac{\pi}{5} y^5 \Big|_0^2 = \frac{32\pi}{5}$

**1997 Calculus AB Solutions: Part A**

24. B Let  $[0,1]$  be divided into 50 subintervals.  $\Delta x = \frac{1}{50}$ ;  $x_1 = \frac{1}{50}, x_2 = \frac{2}{50}, x_3 = \frac{3}{50}, \dots, x_{50} = 1$

Using  $f(x) = \sqrt{x}$ , the right Riemann sum  $\sum_{i=1}^{50} f(x_i)\Delta x$  is an approximation for  $\int_0^1 \sqrt{x} dx$ .

25. A Use the technique of antiderivatives by parts, which was removed from the AB Course Description in 1998.

$$u = x \quad dv = \sin 2x dx$$

$$du = dx \quad v = -\frac{1}{2} \cos 2x$$

$$\int x \sin(2x) dx = -\frac{1}{2}x \cos(2x) + \int \frac{1}{2} \cos(2x) dx = -\frac{1}{2}x \cos(2x) + \frac{1}{4} \sin(2x) + C$$

76. E  $f(x) = \frac{e^{2x}}{2x}$ ;  $f'(x) = \frac{2e^{2x} \cdot 2x - 2e^{2x}}{4x^2} = \frac{e^{2x}(2x-1)}{2x^2}$

77. D  $y = x^3 + 6x^2 + 7x - 2 \cos x$ . Look at the graph of  $y'' = 6x + 12 + 2 \cos x$  in the window  $[-3, -1]$  since that domain contains all the option values.  $y''$  changes sign at  $x = -1.89$ .

78. D  $F(3) - F(0) = \int_0^3 f(x) dx = \int_0^1 f(x) dx + \int_1^3 f(x) dx = 2 + 2.3 = 4.3$

(Count squares for  $\int_0^1 f(x) dx$ )

79. C The stem of the questions means  $f'(2) = 5$ . Thus  $f$  is differentiable at  $x = 2$  and therefore continuous at  $x = 2$ . We know nothing of the continuity of  $f'$ . I and II only.


80. A  $f(x) = 2e^{4x^2}$ ;  $f'(x) = 16xe^{4x^2}$ ; We want  $16xe^{4x^2} = 3$ . Graph the derivative function and the function  $y = 3$ , then find the intersection to get  $x = 0.168$ .

81. A Let  $x$  be the distance of the train from the crossing. Then  $\frac{dx}{dt} = 60$ .

$$S^2 = x^2 + 70^2 \Rightarrow 2S \frac{dS}{dt} = 2x \frac{dx}{dt} \Rightarrow \frac{dS}{dt} = \frac{x}{S} \frac{dx}{dt}$$

After 4 seconds,  $x = 240$  and so  $S = 250$ .

Therefore  $\frac{dS}{dt} = \frac{240}{250}(60) = 57.6$



82. B  $P(x) = 2x^2 - 8x$ ;  $P'(x) = 4x - 8$ ;  $P'$  changes from negative to positive at  $x = 2$ .  $P(2) = -8$

83. C  $\cos x = x$  at  $x = 0.739085$ . Store this in  $A$ .  $\int_0^A (\cos x - x) dx = 0.400$

84. C Cross sections are squares with sides of length  $y$ .

$$\text{Volume} = \int_1^e y^2 dx = \int_1^e \ln x dx = (x \ln x - x) \Big|_1^e = (e \ln e - e) - (0 - 1) = 1$$

85. C Look at the graph of  $f'$  and locate where the  $y$  changes from positive to negative.  $x = 0.91$

86. A  $f(x) = \sqrt{x}$ ;  $f'(x) = \frac{1}{2\sqrt{x}}$ ;  $\frac{1}{2\sqrt{c}} = 2 \cdot \frac{1}{2\sqrt{1}} \Rightarrow c = \frac{1}{4}$

## 1997 Calculus AB Solutions: Part B

87. B  $a(t) = t + \sin t$  and  $v(0) = -2 \Rightarrow v(t) = \frac{1}{2}t^2 - \cos t - 1$ ;  $v(t) = 0$  at  $t = 1.48$
88. E  $f(x) = \int_a^x h(x)dx \Rightarrow f(a) = 0$ , therefore only (A) or (E) are possible. But  $f'(x) = h(x)$  and therefore  $f$  is differentiable at  $x = b$ . This is true for the graph in option (E) but not in option (A) where there appears to be a corner in the graph at  $x = b$ . Also, Since  $h$  is increasing at first, the graph of  $f$  must start out concave up. This is also true in (E) but not (A).
89. B  $T = \frac{1}{2} \cdot \frac{1}{2} (3 + 2 \cdot 3 + 2 \cdot 5 + 2 \cdot 8 + 13) = 12$
90. D
- |                                |  |     |
|--------------------------------|--|-----|
| $F(x) = \frac{1}{2} \sin^2 x$  | $F'(x) = \sin x \cos x$                        | Yes |
| $F(x) = \frac{1}{2} \cos^2 x$  | $F'(x) = -\cos x \sin x$                       | No  |
| $F(x) = -\frac{1}{4} \cos(2x)$ | $F'(x) = \frac{1}{2} \sin(2x) = \sin x \cos x$ | Yes |